

# ***Spitzer/MIPS Photometry of the Extrasolar Planet HD 209458b During Secondary Eclipse***

Drake Deming

*Center for Astrobiology and Planetary Systems Laboratory  
Code 693, NASA's Goddard Space Flight Center  
Greenbelt, MD 20771  
USA  
ddeming@pop600.gsfc.nasa.gov*

Sara Seager

*Department of Terrestrial Magnetism  
Carnegie Institution of Washington  
Washington, DC 20015  
USA*

L. Jeremy Richardson

*National Research Council, and  
Exoplanet and Stellar Astrophysics Branch  
NASA's Goddard Space Flight Center  
Greenbelt MD 20771  
USA*

Joseph Harrington

*Center for Radiophysics and Space Research  
Cornell University  
Ithaca, NY 14853  
USA*

The thermal infrared (IR) flux emitted by a transiting extrasolar planet can in principle be measured by detecting the secondary eclipse, when the planet passes behind the star and re-emerges. For the "hot Jupiter" class of extrasolar planets, the fractional change in the combined IR light of the system during secondary eclipse is predicted from some models to be in the range of 0.2-0.4 percent. This is orders of magnitude greater than the infrared flux expected from both giant extrasolar planets at larger orbital radii, as well as from terrestrial extrasolar planets. The secondary eclipse of the brightest transiting system (HD 209458) was the target of our General Observer program using the MIPS photometer on the Spitzer Space Telescope, at a wavelength of 24 microns. An approximately 6-hour sequence of over 1700 10-second exposures, centered on phase 0.5, was obtained by Spitzer in December 2004, and analysis of these data is currently underway. A preliminary assessment, as of January 2005, indicates that the sensitivity will be limited by the statistical fluctuations in the thermal emission from zodiacal dust in our own solar system, with the limiting sensitivity being sufficient to detect the secondary eclipse if the planet is as hot as predicted by several models ( $> 1200$  Kelvins). We will present our final results, and discuss the prospects for follow-up observations of infrared radiation from transiting extrasolar planets using the James Webb Space Telescope.